



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

∴ (a) would be the better investment.

$120 \times \$13.50 = \$1620$ ;  $\$1620 \times .10\frac{1}{2} \times \frac{1}{12} = \$14.175 =$  profit of first investment over second.

2. (a) Let  $\$13.50 = p$ ,  $.00875 = r$ ,  $\$1000 = a$ ; then  $(a-p) =$  amount due first of first month;  $(a-p)(1+r)-p =$  amount due first of second month;  $[(a-p)(1+r-p)](1+r)-p =$  amount due first of third month;  $\{[(a-p)(1+r-p)](1+r-p)\}(1+r)-p =$  amount due first of fourth month.

∴ for any month, as 10,

$\{[(a-p)(1+r)-p](1+r)-p\}(1+r)-p \} [(a-p)(1+r)-p](1+r)-p^7(1+r)-p$ .

2.  $a(1+r)-p$  at end of first month;  $[a(1+r)-p](1+r)-p$  at end of second month;  $\{[a(1+r)-p](1+r)-p\}(1+r)-p$  at end of third month. At end of tenth month,  $\{ \{ [a(1+r)-p](1+r)-p \}^8(1+r)-p \}$ .

The solution by Dr. Zerr seems to us to be a proper disposition of the problem, but we publish the other solutions for comparison.

Reply to Note of Dr. Drummond on No. 78.—Its one *real* point is, that my solution, on page 43, line 5, omits to denote  $-xy = -6$ , as  $-x_1y_1 = -6$ , (which please correct). Thus result  $x^4 - 8x^2 + 16 = 0$ ;  $y^4 - 18y^2 + 81 = 0$ , whose roots are:  $x=2$  to  $y=3$ ;  $x_a=-2$  to  $y_a=-3$ , of  $xy$  in (I)(II); and also  $x_1=2_1$  to  $y_1=3_1$ ;  $x_2=-2_1$  to  $y_2=-3_1$  (by last two clauses on page 44), the factors of  $-x_1y_1 = -6_1$  in parity to  $x_{11}y_{11} = 35 \dots (D)$  on page 44. Or if the objector insist,—of  $x_1y_1 = 6_1$  and  $x_ay_a = 6_a$  reading then  $xy = 6$  and  $x_ay_a = 6_a$ . No other error is specified or shown. As the equations cited do not occur in my solution,—for his  $x^2 = 36$  is quadratic while  $x^2y^2 = 36$  from (I)(II) is bi-quadratic; I need only say that the roots of his  $x^4 = 1296$  are 6;  $-6$ ,  $6_1/(-1)$ ;  $-6_1/(-1)$ , not  $\pm 6$ ;  $\mp 6$ . Roots  $x^4 = 16$  are 2;  $-2$ ;  $2_1/(-1)$ ;  $-2_1/(-1)$  and not  $\pm 2$ ;  $\mp 2$  as he mistakes me to mean.

J. M. BOORMAN.

## GEOMETRY.

99. Proposed by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio State University, Athens, Ohio.

Find the locus of the vertices of all right cones which have the same given ellipse as a base.

Solution by the PROPOSER.

Any generator,  $x = a'z + \alpha \dots (1)$ .  $y = b'z + \beta \dots (2)$ , must satisfy

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \dots (3), \text{ and } z = 0 \dots (4), \text{ giving } \frac{\alpha^2}{a^2} + \frac{\beta^2}{b^2} = 1 \dots (5).$$

If the vertex be  $(x', y', z')$ , (1) and (2) must be satisfied, and we have  $x' = a'z' + \alpha \dots (6)$ ,  $y' = b'z' + \beta \dots (7)$ .

Eliminating  $a'$ ,  $b'$ ,  $\alpha$ ,  $\beta$  from (1), (2), (6), (7),

$$\frac{1}{a^2}(z'x - x'z)^2 + \frac{1}{b^2}(z'y - y'z)^2 = (z' - z)^2 \dots (8),$$

the equation to a cone having  $(x', y', z')$  for vertex, and (3) for base.

$$(8) \text{ is } \frac{z'^2}{a^2}x^2 + \frac{z'^2}{b^2}y^2 + \left(\frac{x'^2}{a^2} + \frac{y'^2}{b^2} - 1\right)z^2 - \frac{2y'z'}{b^2}yz - \frac{2x'z'}{a^2}xz + 2z'z - z'^2 = 0 \dots (9).$$

The conditions for (9) to be a cone of revolution are  $x'=0$ ,

$$\frac{z'^2}{a^2} - \frac{y'^2}{b^2 - a^2} = 1 \dots (10),$$

an hyperbola for the required locus.

Also solved by G. B. M. ZERR.

100. Proposed by CHARLES CARROLL CROSS, Libertytown, Md.

$O, O_1, O_2, O_3$  are the centers of the inscribed and three escribed circles of a triangle  $ABC$ . Prove  $AO \cdot AO_1 \cdot AO_2 \cdot AO_3 = AB^2 \cdot AC^2$ .

I. Solution by the PROPOSER.

Consider the ex-central triangle  $ABC$  as the original triangle. Then  $H_a H_b H_c$  is the pedal triangle, and the incenter  $O$  becomes the orthocenter  $H$ .

Hence we have to prove  $AH_c \times BH_c \times CH_c \times HH_c = H_a H_c^2 \times H_b H_c^2$ .

We readily find by trigonometry that

$$AH_c = \frac{b^2 + c^2 - a^2}{2c}, \quad BH_c = \frac{a^2 + c^2 - b^2}{2c},$$

$$CH_c = \frac{\sqrt{[4a^2c^2 - (a^2 + c^2 - b^2)^2]}}{2c} = \frac{2\Delta}{c}, \quad HH_c = \frac{(b^2 + c^2 - a^2)(a^2 + c^2 - b^2)}{8c\Delta},$$

$$H_a H_c = \frac{b(a^2 + c^2 - b^2)^2}{2ac}, \quad H_b H_c = \frac{a(b^2 + c^2 - a^2)^2}{2bc}.$$

Substituting in the problem we have

$$\begin{aligned} \frac{b^2 + c^2 - a^2}{2c} \times \frac{a^2 + c^2 - b^2}{2c} \times \frac{2\Delta}{c} \times \frac{(b^2 + c^2 - a^2)(a^2 + c^2 - b^2)}{8c\Delta} \\ = \frac{b^2(a^2 + c^2 - b^2)^2}{4a^2c^2} \times \frac{a^2(b^2 + c^2 - a^2)^2}{4b^2c^2}. \end{aligned}$$

Since these equations cancel, the proposition is proved.

Mr. Cross should have been credited with solutions of problems 96 and 98 in Geometry, and 100 and 101 in Arithmetic.

